

## Sulfur chemistry in the wake of comet Shoemaker-Levy 9

Kevin Zahnle

NASA Ames Research Center, Moffett Field, California

Mordecai-Mark Mac Low

Department of Astronomy and Astrophysics, University of Chicago

Katharina Lodders and Bruce Fegley, Jr.

Department of Earth and Planetary Sciences, Washington University, St. Louis, Missouri

**Abstract.** A curious and unexpected result of the impact of P/Shoemaker Levy 9 with Jupiter was the production of enormous amounts of molecular sulfur ( $S_2$ ). Here we show that  $S_2$  is the natural product of disequilibrium chemistry at low pressures in shocked jovian air, its formation a byproduct of hydrogen recombination. The species observed by the HST —  $S_2$ ,  $CS_2$ , and  $H_2S$  — imply that the G fragment penetrated the  $NH_4SH$  cloud but did not reach the water table. A typical impact within or below the  $NH_4SH$  clouds produces about 0.03-0.1 impactor masses of  $S_2$ . Because comets are relatively hydrogen-poor,  $SO_2$ , not  $S_2$ , is the major product of shocking a water-rich comet, while  $S_2$ ,  $CS_2$ , and  $OCS$  are major products of a desiccated comet. In all cases we find that as the gas cools  $S_2$  converts to the stable low temperature allotrope  $S_8$ , although other chemical fates not modeled here might intervene first.

### Introduction

Diatomic sulfur was observed in absorption by the Hubble Space Telescope in the near UV for the large G impact [Noll *et al.*, 1995]. The observation was made about  $10^4$  seconds after the impact took place, at the center of the impact site, over a footprint roughly 3000 km across. The reported column abundance was  $5 \times 10^{-4}$  g  $cm^{-2}$  [Noll *et al.*, 1995], although this may be an overestimate [R. Yelle, *pers comm.*, 1995]. If the sulfur were uniformly distributed over the entire impact site, a region more than  $10^4$  km across, the mass of  $S_2$  would be of order  $5 \times 10^{14}$  g. Even if restricted to the somewhat smaller region actually observed, the impact produced some  $5 \times 10^{13}$  g of  $S_2$ . These are enormous masses of sulfur. Where did it come from? Why  $S_2$ ? Why so much? Where did it go?

### Model Description

In this paper we address sulfur chemistry in shocked jovian and cometary gas using both an explicitly time-dependent kinetics model and a comprehensive thermochemical equilibrium model, the latter previously developed to study Jupiter at rest [Fegley and Lodders, 1994a]. The kinetics model follows the evolving chemical composition of a parcel of gas by directly integrating the network of chemical reactions. At high temperatures and high pressures chemical reactions are fast and the gas very nearly in thermochemical equilibrium. As the parcel cools and/or rarefies the chemistry increasingly lags behind the evolving conditions, and so becomes increasingly disequibrated. It is often useful to describe the final state as "quenched"; the frozen disequilibrium approximates the equilibrium abundances at the

quench temperature [Zel'dovich and Raizer, 1967, pp. 564-571; Prinn and Fegley, 1987]; Zahnle, 1990].

In an oxygen-poor environment (*i.e.*, dry jovian air) we include 33 chemically active species:  $H_2$ ,  $H$ ,  $C$ ,  $CH$ ,  $CH_2$ ,  $CH_3$ ,  $CH_4$ ,  $C_2$ ,  $C_2H$ ,  $C_2H_2$ ,  $C_2H_3$ ,  $C_2H_4$ ,  $C_2H_5$ ,  $C_2H_6$ ,  $C_4H$ ,  $C_4H_2$ ,  $N$ ,  $NH$ ,  $NH_2$ ,  $NH_3$ ,  $CN$ ,  $HCN$ ,  $N_2$ ,  $S$ ,  $S_2$ ,  $S_3$ ,  $S_4$ ,  $S_8$ ,  $HS$ ,  $H_2S$ ,  $CS$ ,  $CS_2$ ,  $NS$ . Photolysis is neglected. With oxygen this list of species is supplemented by  $O$ ,  $O_2$ ,  $OH$ ,  $H_2O$ ,  $CO$ ,  $CO_2$ ,  $HCO$ ,  $H_2CO$ ,  $NO$ ,  $SO$ ,  $OCS$ ,  $SO_2$ . The latter system is needed to address the chemistry of wet jovian air; it can also be used to approximate the chemistry of cometary material. Reaction rates and sources will be listed elsewhere. Not all the important reactions have measured rates. Unmeasured but plausibly important reactions must be estimated, which leaves considerable scope for error.

The thermochemical equilibrium calculations provide alternative results that are independent of estimated reaction rates. Although the two approaches (kinetic, equilibrium) are not always in complete agreement, they do agree in general, and where they disagree the source of the disagreement can be explained by plausible disequilibrium processes.

### Parcel Histories

We consider two kinds of parcel histories. One kind models the explosion and the subsequent expansion of the fireball. The initial shock conditions, mass of shocked material, and expansion velocity are specified to agree with those generated by a strong point explosion. Thereafter the parcel cools adiabatically as it expands. The second kind models the impact of the plume with the atmosphere. In the latter models the pressure is held constant at a low value (of order 1  $\mu$ bar to 1 mbar), characteristic of pressures in the reentry shock [Zahnle and Mac Low, 1995]. Note that a given parcel is generally shocked twice; *i.e.*, a parcel shocked near the impact site is ejected at high velocity and is shocked again when it reenters the atmosphere. Hence the final state of the gas depends on the second shock. Figure 1 shows an illustrative parcel history taken directly from a numerical model.

**The fireball.** The mass of air  $M_f$  shocked to a temperature greater than  $T$  is approximately

$$25 m_i \left( \frac{2000}{T} \right) < M_f(>T) < 100 m_i \left( \frac{2000}{T} \right). \quad (1)$$

Eq. 1 assumes an impact velocity of 60 km/s, a mean molecular weight of 2.4, and  $\gamma=1.4$ . The lower bound approximates the Sedov solution for a strong explosion and the upper bound assumes that the energy of the impact is spread uniformly over the entire mass of shocked gas; *i.e.*, the upper bound would apply if the shocked gas were well mixed.


**Atmospheric reentry.** The mass of air accelerated to a velocity greater than  $v$  by a typical SL-9 explosion is [Zahnle and Mac Low, 1995]

$$M_r(>v) = 100 m_i \left( \frac{1 \text{ km/s}}{v} \right)^\alpha \quad (2)$$

Copyright 1995 by the American Geophysical Union.

Paper number 95GL01190

0094-8534/95/95GL-01190\$03.00



**Figure 1.** Temperature and pressure history of a representative shocked parcel of jovian air. The parcel originated at the 2 bar level, near the site of the airburst. It was shocked to  $T > 9000$  K, expanded, cooled, and ultimately ejected at about 6 km/sec.

where  $\alpha \approx 1.5$ , and  $v < 15$  km/s. The peak shock temperature is

$$T \geq \frac{\gamma-1}{2} \frac{m}{k} v_z^2 \approx 1500 \text{ K} \left( \frac{\mu}{2.4} \right) \left( \frac{v_z}{5 \text{ km/s}} \right)^2 \quad (3)$$

where the equality is for a strong shock, and  $v_z$  is the normal component of the reentry velocity.  $T'$  can be much higher if the gas is polluted with cometary material, because the cometary material raises the mean molecular weight. Material ejected at 5 km/s falls about  $r \approx v^2/g \approx 1000$  km from the impact site.


We assume a cooling curve in the reentry shock of

$$T(t) = T' \left( \frac{t_0}{t_0 + t} \right)^{1/3} + 120, \quad (4)$$

where  $T'$  denotes the peak shock temperature. We have taken  $t_0 = 20$  s. It must be emphasized that Eq. 4, while generically representative of real parcels, is essentially an arbitrary construct.

## Results

Figure 2 focuses on the sulfur chemistry of a shocked parcel of dry jovian air, as calculated by the kinetics model. The initial chemical composition follows *de Pater* [1991]: compared to solar




**Figure 2.** Time-dependent sulfur chemistry in a representative parcel of shocked dry jovian air. The radicals HS and S are omitted for clarity, but H, which drives the disequilibrium chemistry, is shown. Temperature is also shown. The low pressures, here 20  $\mu$ bars, in the reentry shock strongly favor the disequilibrium production of  $S_2$ .

abundances, carbon is enhanced by a factor of 2.3, nitrogen by 1.5, and sulfur by a factor of 7. There is no water. The temperature and pressure history of the parcel were designed to approximate the history shown in Figure 1.


The primary products of the dry fireball are  $H_2S$  and  $CS_2$ , which are quenched at roughly 1 bar and 2000 K. The high quench temperature and pressure strongly favor the production of  $H_2S$  (*c.f.*, Fig. 3). By contrast the radicals H, S, HS, and CS continue to react. In this example essentially all the CS is converted to  $CS_2$ . We find that  $S_2$  becomes a major product only for peak shock temperatures exceeding about  $10^4$  K, for which the quench pressure of an adiabatically expanded parcel is low.

The reentry shock in Fig 2 assumes  $T=2500$  and a pressure of 10  $\mu$  bar. In this particular simulation  $S_2$  is the major sulfur species from 500 to  $2 \times 10^4$ s. As the parcel cools the bulk of the sulfur converts to the stable low temperature allotrope  $S_8$ . The period of  $S_2$  dominance is controlled by the cooling timescale and the pressure. At 1 mbar conversion to  $S_8$  occurs at 500 K, at 10  $\mu$ bar conversion is delayed until T drops to 250 K. At low temperatures the thermodynamically preferred  $H_2S$  reappears when atomic hydrogen disappears.

Production of  $S_2$  in a cooling hydrogen-rich gas is driven by the recombination of molecular hydrogen. The key reactions are



All three reactions are exothermic. The net reaction is



The equilibrium increasingly favors the products as the temperature drops.

We calculate chemical products of the fireball by dividing the shocked air into concentric shells. Each shell is shocked to a peak temperature  $T'$ . The mass of a shell is determined from Eq. 1. The final chemical products are summed over all shells. Chemical products for wet and dry air are listed in Table 1 for a  $10^{14}$  g impactor; the numbers given in the Table are based on the upper bound from Eq. 1. If the lower bound is adopted yields drop by a factor of ten.

Eqs. 2 and 3 determine the mass of air shocked to temperatures greater than  $T'$  in the reentry shock. Each annulus cools from  $T'$  as described by Eq. 4. A constant pressure of 1 mbar was assumed unless otherwise noted; results differ somewhat for other pressures. Final chemical products for a  $10^{14}$  g impactor are listed in Table 2. Essentially all the  $S_8$  in Table 2 was at one time  $S_2$ . Evidently the HST first observed  $S_2$  before it converted to  $S_8$ .

**Table 1.** Sulfur Products from the Fireball (g)<sup>a</sup>

species	WC <sup>b</sup>	DC <sup>c</sup>	DA <sup>d</sup>	WA <sup>e</sup>
$H_2S$ <sup>f</sup>	—	3(11)	>4(13)	>4(13)
$S_2$	2(5)	2(12)	7(12)	6(12)
CS	—	2(5)	1(13)	4(4)
$CS_2$	—	2(12)	8(11)	3(8)
OCS	—	1(12)	—	1(11)
SO	5(8)	2(5)	—	1(10)
SO <sub>2</sub>	5(12)	3(4)	—	2(10)
CO	5(13)	7(13)	—	8(13)
$H_2O$ <sup>f</sup>	4(13)	1(9)	—	>3(14)

<sup>a</sup> Assumes a  $10^{14}$  g comet. 3(11) =  $3 \times 10^{11}$

<sup>b</sup> Wet Comet; solar C:N:S:H<sub>2</sub>O.

<sup>c</sup> Dry Comet; solar C:N:S; H<sub>2</sub>O is 30% solar.

<sup>d</sup> Dry Air; C:N:S enhanced by 2.3:1.5:7 over solar.

<sup>e</sup> Wet Air; assumes solar C/O.

<sup>f</sup> Present in unprocessed air; lower limits only.

## The Comet

The comet provides a second chemical environment, one that is unlike jovian air. We consider two "comet" compositions: a "wet" comet with a solar water to carbon ratio, and a "dry" comet from which 70% of the water has been lost. By construction the comets contain only the hydrogen in water and organic matter, the latter with an assumed C/H ratio of one. Cometary products of the fireball and the reentry shock are listed in Tables 1 and 2.

The reported detection of a great deal of hot water [Bjoraker *pers. comm.* 1994] in the minutes immediately following the impact argues that the comet was water-rich (C/O < 1). According to Table 2 the major sulfur species produced from such a comet would be SO<sub>2</sub>, which, as noted earlier, was not detected by HST. On the other hand, the reported detection of a great deal of OCS [Lellouche *et al.*, 1995] argues that the comet was dry. Here remains a puzzle.


## Thermochemical Equilibrium

Thermochemical equilibrium calculations in solar, dry solar, and cometary gas provide a complementary perspective. We use the CONDOR code and database described by Fegley and Lodders [1994a]. Approximately 2000 gases of all naturally occurring elements in the periodic table were included in these calculations [Fegley and Lodders, 1994b].

At high temperatures (>3500 K, essentially independent of total pressure), ionized and atomic sulfur are the dominant species. With cooling of the fireball, the radicals HS and CS become important, their abundances eventually drop with continued cooling and H<sub>2</sub>S, S<sub>2</sub>, and CS<sub>2</sub> become important at low temperatures. Figure 3 illustrates that the abundances of S<sub>2</sub> and CS<sub>2</sub> are sensitive to temperature and pressure, and both peak at low temperatures (≈1000 K) and low pressures (1 μbar and below). The CS radical constitutes a significant fraction of total sulfur over wide T and P ranges, but is probably unquenchable save at very low pressure. The plausible major sulfur species resulting from shock induced chemical reprocessing of jovian gas are H<sub>2</sub>S, S<sub>2</sub>, CS<sub>2</sub> and at low pressures, CS.

Thermochemical equilibrium calculations show that H<sub>2</sub>S and S<sub>2</sub> also result from shock heating of solar gas, i.e. wet jovian air. But production of CS and CS<sub>2</sub> is suppressed. In their place we find oxides such as SO, SO<sub>2</sub>, and OCS. Shock heating of cometary gas, using compositions for P/Halley given by Anders and Grevesse [1989], also yields H<sub>2</sub>S and S<sub>2</sub>, but again large amounts of SO<sub>2</sub>, SO, and other oxidized sulfur species also form.

Although these preliminary results of the thermochemical equilibrium calculations differ in detail from those of the kinetic model, both sets of calculations show the same major conclusions, namely that S<sub>2</sub> and CS<sub>2</sub> production is a natural conse-



**Figure 3.** Thermochemical equilibria of S<sub>2</sub>, CS<sub>2</sub>, and CS as functions of temperature and pressure in dry solar gas with a C/O atomic ratio of 100. This ratio is equivalent to the CH<sub>4</sub>/H<sub>2</sub>O ratio at 200 K, in the NH<sub>4</sub>SH cloud layer on Jupiter. Isobars from 1 to 10<sup>-9</sup> bars and temperatures from 500 K to 4500 K are considered. Higher temperatures lead to increasing amounts of atomic and ionized sulfur and are not graphed.

**Table 2.** Sulfur Products from the Reentry Shock (g)<sup>g</sup>

species	WC <sup>b</sup>	DC <sup>c</sup>	DA <sup>d</sup>	DA <sup>dh</sup>	WA <sup>e</sup>
H <sub>2</sub> S <sup>f</sup>	5(6)	7(10)	>1(13)	>1(13)	>4(12)
S <sub>2</sub>	2(5)	1(9)	3(9)	5(8)	2(10)
S <sub>8</sub>	—	2(12)	6(12)	4(12)	9(12)
CS	—	—	—	7(12)	—
CS <sub>2</sub>	—	2(12)	4(10)	3(11)	4(8)
OCS	7(7)	4(11)	—	—	5(9)
SO	3(8)	—	—	—	1(11)
SO <sub>2</sub>	5(12)	—	—	—	1(12)
CO	5(13)	7(13)	—	—	5(13)
H <sub>2</sub> O <sup>f</sup>	4(13)	1(7)	—	—	>9(13)

<sup>g</sup> Assumes a 10<sup>14</sup> g comet and p' = 1 mbar.

<sup>h</sup> p' = 0.1 mbar.

quence of shock heating and quenching dry jovian gas. Both sets of calculations also show that shock heating of wet jovian gas and of cometary gas leads to the production of oxidized sulfur gases, exemplified by SO<sub>2</sub> and SO.

## Summary

Diatomic sulfur was a major unexpected product of adding P/Shoemaker-Levy 9 to Jupiter. The S<sub>2</sub> and CS<sub>2</sub> were apparently concentrated at high altitudes, the latter above a millibar [Yelle and McGrath, *pers. comm.*, 1995], which is consistent with the sulfur falling from above; i.e., S<sub>2</sub> and CS<sub>2</sub> were produced where

the hypersonic ejecta plume fell back on the atmosphere. A sulfur content of 0.3% sulfur by mass [de Pater, 1991] would place the reported column [Noll et al., 1995] of S<sub>2</sub> above 400 μbars. Kinetic and thermochemical equilibrium models presented here show that the low pressure and extreme disequilibrium chemistry of the reentry shock strongly favor the generation of S<sub>2</sub>. Eventually S<sub>2</sub> converts to S<sub>8</sub> when the gas cools. Survival of S<sub>2</sub> is favored by low pressure and high temperature. Moses et al. [1995] show that conversion to S<sub>8</sub> is slow above 100 μbars, where low density and photolysis prevent polymerization.

Shocks in dry jovian air produce mostly S<sub>2</sub>, H<sub>2</sub>S, CS<sub>2</sub>, and at low pressures, CS. These were the observed species, and the calculated S<sub>2</sub>/CS<sub>2</sub> ratio is in reasonable agreement with that observed. Shocks in wet jovian air also produce abundant S<sub>2</sub>, but the oxides SO<sub>2</sub> and SO replace CS and CS<sub>2</sub> as important other species. Oxides were not observed by the HST [Noll et al., 1995]. Apparently the comet did not excavate much air from below the jovian water table.

Shocks in the comet itself produce SO<sub>2</sub> if the comet is wet, and CS<sub>2</sub>, S<sub>2</sub>, and OCS if the comet is dry. The detection of hot water implies that the comet was wet, but the detection of OCS implies that the comet was dry. Any comet produces CO. The apparent absence of SO<sub>2</sub> might be explained by spreading the ex-comet over the full breadth of the ejecta blanket. An alternative explanation is that the comet, being relatively dense, is mostly buried beneath the center of the explosion [Crawford et al., 1994]. The latter possibility is consistent with small amounts of SO<sub>2</sub> but inconsistent with large amounts of water, CO, and OCS.

The calculated mass of S<sub>2</sub> produced by a 10<sup>14</sup> g impact falls about an order of magnitude short of the reported 5x10<sup>13</sup> g [Noll et al., 1995]. A 10<sup>15</sup> g impactor is needed to produce this much S<sub>2</sub>. Either Jupiter is considerably more rich in sulfur than we have assumed; the observed amount of S<sub>2</sub> has been overestimated; photolysis by UV radiation from the fireball was much more effective than shocks at driving chemistry; or the impactors were larger than the 10<sup>14</sup> g fragments deduced by Asphaug and Benz [1994] from the gravitational dynamics of tidal disruption and re-accretion and by Zahnle and Mac Low [1995] from the infrared luminosity of the R fragment plume-fall.

**Acknowledgments.** We thank G. Bjoraker, C. P. McKay, K. Noll, I. de Pater, R. Yelle, & J. I. Moses. B.F. & K.L. were supported by NASA

Grant NAGW-2861 from the Planetary Atmospheres program. M.-M. M. L. was supported by NSF Grant AST93-22509 from the Aeronomy program. K.Z. was supported by the NASA Exobiology program.

## References

- Asphaug, E., and Benz, W., Density of comet Shoemaker-Levy 9 deduced by modelling breakup of the parent 'rubble pile', *Nature*, 370, 120-124, 1994.
- Anders, E. and Grevesse, N., Abundances of the elements: meteoritic and solar, *Geochim. Cosmochim. Acta*, 53, 197-214, 1989.
- Fegley, B. and Lodders, K., Chemical models of the deep atmospheres of Jupiter and Saturn, *Icarus*, 110, 117-154, 1994a.
- Fegley, B. and Lodders, K., Chemical consequences of the impact of comet SL9 on Jupiter, *EOS Trans AGU*, 76, 405, 1994b.
- Lellouche, E. et al., Chemical and thermal response of Jupiter to the impact of comet Shoemaker-Levy 9, *Nature*, 373, 592-595, 1995.
- Mac Low, M.-M. and Zahnle, K., Explosion of Comet Shoemaker-Levy 9 on entry into the Jovian atmosphere, *Astrophys. J. Lett.*, 434, L33-L36, 1994.
- Moses J. I., Allen M., and Gladstone R., The photochemical evolution of sulfur compounds following the Shoemaker-Levy 9 impacts with Jupiter, *Geophys. Res. Lett.*, 1995.
- Noll, K. S. et al., Hubble Space Telescope spectroscopic observations of Jupiter after the collision of Comet P/Shoemaker-Levy 9, *Science*, 267, 1307-1313, 1995.
- de Pater, I., The Significance of Radio Observations for Planets, *Physics Reports*, 200, 1-37, 1991.
- Prinn, R. and Fegley, B., Bolide Impacts, acid rain, and biospheric traumas at the Cretaceous-Tertiary boundary, *Earth Planet. Sci. Lett.*, 83, 1-15, 1987.
- Zahnle, K., Atmospheric chemistry by large impacts, *Global Catastrophes in Earth History*, edited by V. Sharpton and P. Ward, G.S.A. Special Paper 247, 271-288, 1990.
- Zahnle, K. and Mac Low, M.-M., The collision of Jupiter and Comet Shoemaker-Levy 9, *Icarus*, 108, 1-17, 1994.
- Zahnle, K. and Mac Low, M.-M., A simple model of the light curve generated by a Shoemaker-Levy 9 impact, *J. Geophys. Res. (submitted)* 1995.

K. Zahnle, NASA Ames Research Center, MS 245--3, Moffett Field, CA 94035-1000

M.-M. Mac Low, Department of Astronomy and Astrophysics, 5640 S. Ellis Ave., University of Chicago, Chicago IL 60637

K. Lodders and B. Fegley, Jr., Department of Earth and Planetary Sciences, Campus Box 1169, Washington University, St. Louis MO 63130-4899

(received December 19, 1994; revised March 10, 1995; accepted March 15, 1995)